## Boyle's Law

1) If I have 5.6 liters of gas in a piston at a pressure of 1.5 atm and compress the gas until its volume is 4.8 L, what will the new pressure inside the piston be?

2) I have added 15 L of air to a balloon at sea level (1.0 atm). If I take the balloon with me to Denver, where the air pressure is 0.85 atm, what will the new volume of the balloon be?

3) I've got a car with an internal volume of 12,000 L. If I drive my car into the river and it implodes, what will be the volume of the gas when the pressure goes from 1.0 atm to 1.4 atm?

## Boyle's Law - Solutions

1) If I have 5.6 liters of gas in a piston at a pressure of 1.5 atm and compress the gas until its volume is 4.8 L, what will the new pressure inside the piston be?

 $P_1V_1 = P_2V_2$ (1.5 atm)(5.6 L) = (x)(4.8 L) x = 1.8 atm

2) I have added 15 L of air to a balloon at sea level (1.0 atm). If I take the balloon with me to Denver, where the air pressure is 0.85 atm, what will the new volume of the balloon be?

 $P_1V_1 = P_2V_2$ (1.0 atm)(15 L) = (0.85 atm)(x) x = 18 L

3) I've got a car with an internal volume of 12,000 L. If I drive my car into the river and it implodes, what will be the volume of the gas when the pressure goes from 1.0 atm to 1.4 atm?  $P_1V_1 = P_2V_2$ (1.0 atm)(12,000 L) = (1.4 atm)(x)